



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12N 15/54, 15/82, A01H 5/00</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/66749</b> <b>(43) International Publication Date:</b> 9 November 2000 (09.11.00)
<b>(21) International Application Number:</b> PCT/US00/12316 <b>(22) International Filing Date:</b> 4 May 2000 (04.05.00) <b>(30) Priority Data:</b> 60/132,417 4 May 1999 (04.05.99) US <b>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application</b> US 60/132,417 (CON) Filed on 4 May 1999 (04.05.99) <b>(71) Applicant (for all designated States except US):</b> CARGILL, INCORPORATED [US/US]; 15407 McGinty Road West, Wayzata, MN 55391 (US). <b>(72) Inventor; and</b> <b>(75) Inventor/Applicant (for US only):</b> SHORROSH, Basil, S. [US/US]; 3311 Laredo Lane, Fort Collins, CO 80526 (US). <b>(74) Agent:</b> ELLINGER, Mark, S.; Fish & Richardson P.C., P.A., 60 South Sixth Street, Suite 3300, Minneapolis, MN 55402 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> PLANT ACYLTRANSFERASES		
<b>(57) Abstract</b>		
<p>A gene encoding diacylglycerol acyltransferase from <i>Brassica napus</i> is described, as well as methods of altering oil content in plants using the diacylglycerol acyltransferase.</p>		

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## PLANT ACYLTRANSFERASES

### TECHNICAL FIELD

The invention relates to alteration of oil content in plants.

### BACKGROUND

5        Triacylglycerols produced in plant tissues (e.g., canola seeds) are a significant source of fatty acids in the human diet. Diets rich in animal fat appear to promote maladies such as heart disease in humans. The ability to modulate triacylglycerol synthesis in plants could allow for production of fatty acid compositions that are more beneficial for humans or more efficiently synthesized by the plant. Cloning and  
10        characterization of plant genes encoding enzymes involved in triacylglycerol synthesis and metabolism represents a major advance toward controlling triacylglycerol synthesis in plants.

### SUMMARY

      A new plant acyltransferase gene from *Brassica napus* has been cloned and  
15        characterized. The protein encoded by this gene is involved in triacylglycerol synthesis. Partial DNA sequences of the *B. napus* gene confirm that it is related to an *Arabidopsis thaliana* putative acyltransferase gene. Antisense expression of the *B. napus* sequence decreases total oil content.

      In one aspect, the invention features an isolated nucleic acid having at least  
20        80% (e.g., 85%, 90%, or 99%) sequence identity to the nucleotide sequence of SEQ ID NO:3, or to a fragment of the nucleotide sequence of SEQ ID NO:3, the fragment being at least 15 nucleotides in length. An isolated nucleic acid having at least 80% (e.g., 85%, 90%, or 99%) sequence identity to the nucleotide sequence of SEQ ID NO:4, or to a fragment of the nucleotide sequence of SEQ ID NO:4, the fragment  
25        being at least 15 nucleotides in length is also featured.

      In another aspect, the invention features an isolated nucleic acid that includes a first and second region, the first region having at least 80% sequence identity to the nucleotide sequence of SEQ ID NO:3 and the second region having at least 80% sequence identity to the nucleotide sequence of SEQ ID NO:4. The nucleic acid can  
30        encode a diacylglycerol acyltransferase polypeptide. The first and second regions can

be separated by approximately 600 nucleotides. The nucleic acid can be the insert of pMB143. An expression vector including the nucleic acid operably linked to an expression control element also is featured. The nucleic acid can be operably linked in antisense orientation.

5           In another aspect, the invention features a transgenic plant and progeny thereof that include an exogenous nucleic acid encoding a diacylglycerol acyltransferase polypeptide operably linked to a regulatory element, and seed produced by such plants. The nucleic acid can include a first and second region, the first region having at least 80% sequence identity to the nucleotide sequence of SEQ ID NO:3, the  
10       second region having at least 80% sequence identity to the nucleotide sequence of SEQ ID NO:4. The transgenic plant can produce seeds that exhibit a statistically significantly altered oil content as compared to seeds produced by a corresponding plant lacking the nucleic acid encoding the diacylglycerol acyltransferase polypeptide. The plant can be a soybean plant or a *Brassica* plant.

15           The invention also features a nucleic acid that includes the nucleotide sequence of SEQ ID NO:3 or SEQ ID NO:4 or the nucleotide sequence exactly complementary to SEQ ID NO:3 or SEQ ID NO:4. The nucleic acid also can include the RNA equivalent of the nucleotide sequence of SEQ ID NO:3 or SEQ ID NO:4, or an RNA equivalent that is exactly complementary to the nucleotide sequence of SEQ  
20       ID NO:3 or SEQ ID NO:4. The nucleotide sequences of SEQ ID NO:3 and SEQ ID NO:4 are partial sequences of the new *B. napus* gene.

          In another aspect, the invention features a nucleic acid that (1) hybridizes under stringent conditions to a DNA molecule consisting of the nucleotide sequence of SEQ ID NO:3 or SEQ ID NO:4, and (2) encodes a plant acyltransferase. In  
25       addition, the invention includes a nucleic acid (1) having a nucleotide sequence which is at least 80% (e.g., at least 82, 85, 90, 92, 95, 98, or 99%) identical to the nucleotide sequence of SEQ ID NO:3 or SEQ ID NO:4, and (2) encoding a plant acyltransferase. As used herein, the term "stringent conditions" means hybridization at 42°C in the presence of 50% formamide; a first wash at 65°C with 2X SSC containing 1% SDS;  
30       followed by a second wash at 65°C with 0.1X SSC.

The nucleotide sequence fragments described below can be used to hybridize against cDNA or genomic DNA libraries from a variety of sources to clone genes related to the *B. napus* acyltransferase gene. In addition, the sequence fragments can be used to design additional primers for further sequencing of the *B. napus* gene or for  
5 PCR amplification of portions of the *B. napus* gene.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used to practice the invention, suitable methods and  
10 materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

15 Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

#### DESCRIPTION OF DRAWINGS

Figure 1 is a schematic of a construct containing the acyltransferase. Restriction sites are marked.

20 Figure 2 is a schematic of a construct containing the acyltransferase in antisense orientation. Restriction sites are marked.

#### DETAILED DESCRIPTION

Triacylglycerols (TAGs) are synthesized by the sequential transfer of acyl chains to the glycerol backbone by a series of enzymes in the Kennedy pathway  
25 (glycerol-3-phosphate acyltransferase, lysophosphatidic acid acyltransferase, and diacylglycerol acyltransferase). See, Töpfer et al., Science, 1995, 268:681-686. Diacylglycerol can be used to form TAGs or membrane glycerolipids, and is the substrate for diacylglycerol acyltransferase (DAGAT, E.C. 2.3.1.20). DAGAT transfers a third acyl chain to diacylglycerol, forming a TAG, and is the only enzyme  
30 unique to TAG synthesis in the Kennedy pathway. Thus, the reaction catalyzed by

DAGAT represents a key branchpoint in TAG synthesis. As described herein, a 1.4 kB gene that encodes DAGAT has been identified from *Brassica* and can be used to alter total oil content in plants.

*DAGAT Nucleic Acid Molecules*

5           The invention features isolated nucleic acids having at least 80% sequence identity, e.g., 85%, 90%, 95%, or 99% sequence identity, to the nucleic acid of SEQ ID NO:3 or SEQ ID NO:4, or fragments of the nucleic acid of SEQ ID NO:3 or 4 that are at least about 15 nucleotides (nt) in length (e.g., at least 18, 20, 22, 24, 26, 28, or 30 nt). In one embodiment, the nucleic acid includes a first region having at least  
10   80% sequence identity to the nucleotide sequence of SEQ ID NO:3 and a second region having at least 80% sequence identity to the nucleotide sequence of SEQ ID NO:4. The first and second regions can be separated by about 590 to about 700 nucleotides, e.g., about 600 nucleotides.

          Generally, percent sequence identity is calculated by determining the number  
15   of matched positions in aligned nucleic acid sequences, dividing the number of matched positions by the total number of aligned nucleotides, and multiplying by 100. A matched position refers to a position in which identical nucleotides occur at the same position in aligned nucleic acid sequences. The total number of aligned nucleotides refers to the number of nucleotides from SEQ ID NO:3 or NO:4 that are  
20   being aligned. Nucleic acid sequences can be aligned by the Clustal algorithm of MEGALIGN® (DNASTAR, Madison, WI, 1997) sequence alignment software. In this method, sequences are grouped into clusters by examining the distances between all pairs. Clusters are aligned as pairs, then as groups. A gap penalty of 100 and a gap length penalty of 2 are used in the alignments.

25           Isolated nucleic acid molecules of the invention can be produced by standard techniques. As used herein, "isolated" refers to a sequence corresponding to part or all of a gene encoding a DAGAT polypeptide, but free of sequences that normally flank one or both sides of the wild-type gene in a plant genome. As used herein, "polypeptide" refers to a chain of least eight amino acids. An isolated nucleic acid  
30   can be, for example, a DNA molecule, provided one of the nucleic acid sequences normally found immediately flanking that DNA molecule in a naturally-occurring

genome is removed or absent. Thus, an isolated nucleic acid includes, without limitation, a DNA molecule that exists as a separate molecule (e.g., a cDNA or genomic DNA fragment produced by PCR or restriction endonuclease treatment) independent of other sequences as well as recombinant DNA that is incorporated into  
5 a vector, an autonomously replicating plasmid, a virus (e.g., a retrovirus, adenovirus, or herpes virus), or into the genomic DNA of a prokaryote or eukaryote. In addition, an isolated nucleic acid can include a recombinant DNA molecule that is part of a hybrid or fusion nucleic acid. A nucleic acid existing among hundreds to millions of other nucleic acids within, for example, cDNA libraries or genomic libraries, or gel  
10 slices containing a genomic DNA restriction digest, is not to be considered an isolated nucleic acid.

Isolated nucleic acid molecules are at least about 15 nucleotides in length. For example, the nucleic acid molecule can be about 20 to 35, 40-50, 50-100, or greater than 150 nucleotides in length, e.g., 200-300, 300-500, 500-1000, or 1100-1500  
15 nucleotides in length. Such fragments, whether encoding a polypeptide or not, can be used as probes, primers, and diagnostic reagents. In some embodiments, the isolated nucleic acid molecules encode a full-length DAGAT polypeptide. Nucleic acid molecules can be DNA or RNA, linear or circular, and in sense or antisense orientation. The nucleic acid molecules also can be complementary to the nucleotide  
20 sequences of SEQ ID NO:3 or SEQ ID NO:4. A nucleic acid encoding a DAGAT polypeptide may or may not contain introns within the coding sequence.

Polymerase chain reaction (PCR) techniques can be used to produce nucleic acid molecules of the invention. PCR refers to a procedure or technique in which target nucleic acids are amplified. Sequence information from the ends of the region  
25 of interest or beyond typically is employed to design oligonucleotide primers that are identical in sequence to opposite strands of the template to be amplified. PCR can be used to amplify specific sequences from DNA as well as RNA, including sequences from total genomic DNA or total cellular RNA. Primers are typically 14 to 40 nucleotides in length, but can range from 10 nucleotides to hundreds of nucleotides in  
30 length. General PCR techniques are described, for example in PCR Primer: A Laboratory Manual, Ed. by Dieffenbach, C. and Dveksler, G., Cold Spring Harbor Laboratory Press, 1995.

Nucleic acids encoding DAGAT polypeptides also can be produced by chemical synthesis, either as a single nucleic acid molecule or as a series of oligonucleotides. For example, one or more pairs of long oligonucleotides (e.g., >100 nucleotides) can be synthesized that contain the desired sequence, with each pair  
5 containing a short segment of complementarity (e.g., about 15 nucleotides) such that a duplex is formed when the oligonucleotide pair is annealed. DNA polymerase is used to extend the oligonucleotides, resulting in a double-stranded nucleic acid molecule per oligonucleotide pair, which then can be ligated into a vector.

It should be appreciated that many different nucleic acids will encode a  
10 polypeptide having a particular DAGAT amino acid sequence. The degeneracy of the genetic code is well known in the art, *i.e.*, many amino acids are encoded by more than one nucleotide codon. It should also be appreciated that certain amino acid substitutions can be made within polypeptide sequences without affecting the function of the polypeptide. Conservative amino acid substitutions or substitutions of similar  
15 amino acids often are tolerated without affecting polypeptide function. Similar amino acids can be those that are similar in size and/or charge properties. Similarity between amino acids has been assessed in the art. For example, Dayhoff et al. (1978) in *Atlas of Protein Sequence and Structure*, Vol. 5, Suppl. 3, pp. 345-352, provides frequency tables for amino acid substitutions that can be employed as a measure of amino acid  
20 similarity.

#### *Transgenic Plants*

The invention features transgenic plants that have altered total oil content, *i.e.*, increased or decreased oil content. Suitable plant species include, for example, *Brassica* spp. such as *B. napus*, *B. campestris*, *B. juncea*, and *B. rapa* (canola-type  
25 and high erucic acid rapeseed), soybean, sunflower, castor bean, safflower, crambe, palm, coconut, corn, cottonseed, olive, peanut, flax, and sesame. Canola, soy, sunflower, and safflower plants having increased oil content are particularly useful. Table 1 provides relative percent oil and protein content on a dry weight basis (unless indicated otherwise) of suitable oilseed plants.

30 The present invention describes a novel method of making transgenic plants that produce seeds with a statistically significant alteration in oil content. As used



herein, "statistically significant" refers to a p-value of less than 0.05, *e.g.*, a p-value of less than 0.025 or a p-value of less than 0.01, using an appropriate measure of statistical significance, *e.g.*, a one-tailed two sample t-test. By using this method, plants can be produced that exhibit an altered oil content in their seeds. The altered oil content is statistically significant relative to the oil content of unmodified seeds. Plants produced by the method of the present invention can produce seeds having an increase in oil of from about 1% to about 25% over the oil content in seeds produced by unmodified control plants. For example, the increase in oil content for plants described herein can be from about 2% to about 20%, from about 4% to about 15%, from about 5% to about 10%, or from about 10% to about 20%, relative to unmodified plants.

**TABLE 1**  
**Relative Percent Oil and Protein Content of Oilseed Plants**

Plant	% Oil	% Protein	Key
Soybean ( <i>Glycine max</i> )	~20	~40	C
Rapeseed ( <i>Brassica napus</i> )	40-44	38-41 (oil free meal)	C; D
Sunflower ( <i>Helianthus annus</i> )	40		D
Castor bean ( <i>Ricinus communis</i> )	50		A
Safflower ( <i>Carthamus tinctorius</i> )	36.8-47.7	15.4-22.5	D
Crambe ( <i>Crambe abyssinica</i> )	30-35	~28	B
Palm ( <i>Elaeis guineensis</i> )	20 >50		C; per fresh fruit bunch (~20% moisture); Dried kernels
Coconut ( <i>Cocos nucifera</i> )	34 69	3.5	D; coconut flesh (50% moisture); dried kernels
Maize ( <i>Zea mays</i> )	3.1-5.7	6-12	C; D
Cottonseed ( <i>Gossypium hirsutum</i> )	25-30	25-30	D; kernel
Olive ( <i>Olea europaea</i> )	19.6	1.6	fruit (52.4% moisture)
Peanut ( <i>Arachis hypogaea</i> )	36-56	25-30	C; (unknown moisture)
Flax ( <i>Linum usitatissimum</i> )	35-45		D; per fruit capsule (~10 seeds/fruit)

Sesame ( <i>Sesamum indicum</i> )	53.3-57.5	25-30	D; (5-7% moisture)
--------------------------------------	-----------	-------	--------------------

(A) Brigham RD, 1993, Castor: Return of an old crop, p 380-3. In *New Crops*, Janick, J & Simon, JE, eds. Wiley, NY.

(B) Grombacher et al., Cooperative Extension, Institute of Agriculture and Natural Resources, University of Nebraska-Lincoln, *Crambe production*, Publication G93-1126A, G1126 (Field Crops), F-17 (Misc. Crops); see also [pubs@unlvm.unl.edu](mailto:pubs@unlvm.unl.edu)

(C) In *Principles of Cultivar Development*, 1987, Fehr, WR, ed., Macmillan Publishing Co., NY.

(D) In *5<sup>th</sup> Edition Bailey's Industrial Oil & Fat Products*, Vol. 2, Edible Oil & Fat Products: Oils and Oil Seeds, 1996, Hui, YH, ed., Wiley, NY.

A plant described herein may be used as a parent to develop a plant line, or may itself be a member of a plant line, i.e., it is one of a group of plants that display little or no genetic variation between individuals for total oil content. Such lines can be created by several generations of self-pollination and selection, or vegetative propagation from a single parent using tissue or cell culture techniques known in the art. Additional means of breeding plant lines from a parent plant are known in the art.

In general, plants of the invention can be obtained by introducing at least one exogenous nucleic acid encoding a DAGAT polypeptide into plant cells. As used herein, the term "exogenous" refers to a nucleic acid that is introduced into the plant. Exogenous nucleic acids include those that naturally occur in the plant and have been introduced to provide one or more additional copies, as well as nucleic acids that do not naturally occur in the plant. Typically, a nucleic acid construct containing a nucleic acid encoding a DAGAT polypeptide is introduced into a plant cell. Seeds produced by a transgenic plant can be grown and selfed (or outcrossed and selfed) to obtain plants homozygous for the construct. Seeds can be analyzed to identify those homozygotes having the desired expression of the construct. Transgenic plants can be entered into a breeding program, e.g., to increase seed, to introgress the novel construct into other lines or species, or for further selection of other desirable traits. Alternatively, transgenic plants can be obtained by vegetative propagation of a transformed plant cell, for those species amenable to such techniques.

Progeny of a transgenic plant are included within the scope of the invention, provided that such progeny exhibit altered oil content. Progeny of an instant plant

include, for example, seeds formed on F<sub>1</sub>, F<sub>2</sub>, F<sub>3</sub>, and subsequent generation plants, or seeds formed on BC<sub>1</sub>, BC<sub>2</sub>, BC<sub>3</sub>, and subsequent generation plants.

Transgenic techniques for use in the invention include, without limitation, *Agrobacterium*-mediated transformation, electroporation, and particle gun transformation. Illustrative examples of transformation techniques are described in WO 99/43202 and U.S. Patent 5,204,253 (particle gun) and U.S. Patent 5,188,958 (*Agrobacterium*). Transformation methods utilizing the Ti and Ri plasmids of *Agrobacterium* spp. typically use binary type vectors. Walkerpeach, C. et al., in Plant Molecular Biology Manual, S. Gelvin and R. Schilperoort, eds., Kluwer Dordrecht, C1:1-19 (1994). If cell or tissue cultures are used as the recipient tissue for transformation, plants can be regenerated from transformed cultures by techniques known to those skilled in the art. In addition, various plant species can be transformed using the pollen tube pathway technique.

Nucleic acid constructs suitable for producing transgenic plants of the invention include a nucleic acid encoding a DAGAT polypeptide operably linked to a regulatory element such as a promoter. Standard molecular biology techniques can be used to generate nucleic acid constructs. To increase oil content in plants, the nucleic acid encoding a DAGAT polypeptide is operably linked to the regulatory element in sense orientation.

Suitable promoters can be constitutive or inducible, and can be seed-specific. As used herein, "constitutive promoter" refers to a promoter that facilitates the expression of a nucleic acid molecule without significant tissue- or temporal-specificity. An inducible promoter may be considered to be a "constitutive promoter", provided that once induced, expression of the nucleic acid molecule is relatively constant or uniform without significant tissue- or temporal-specificity. Suitable promoters are known (*e.g.*, Weising *et al.*, *Ann. Rev. Genetics* 22:421-478 (1988)). The following are representative examples of promoters suitable for use herein: regulatory sequences from fatty acid desaturase genes (*e.g.*, *Brassica fad2D* or *fad2F*, see WO 00/07430); alcohol dehydrogenase promoter from corn; light inducible promoters such as the ribulose biphosphate carboxylase (Rubisco) small subunit gene promoters from a variety of species; major chlorophyll a/b binding

protein gene promoters; the 19S promoter of cauliflower mosaic virus (CaMV); as well as synthetic or other natural promoters that are either inducible or constitutive. In one embodiment, regulatory sequences are seed-specific, i.e., the particular gene product is preferentially expressed in developing seeds and expressed at low levels or not at all in the remaining tissues of the plant. Non-limiting examples of seed-specific promoters include napin, phaseolin, oleosin, and cruciferin promoters.

Additional regulatory elements may be useful in the nucleic acid constructs of the present invention, including, but not limited to, polyadenylation sequences, enhancers, introns, and the like. Such elements may not be necessary for expression of a DAGAT polypeptide, although they may increase expression by affecting transcription, stability of the mRNA, translational efficiency, or the like. Such elements can be included in a nucleic acid construct as desired to obtain optimal expression of the acyltransferase nucleic acid in the host cell(s). Sufficient expression, however, may sometimes be obtained without such additional elements. A reference describing specific regulatory elements is Weising *et al.*, *Ann. Rev. Genetics* 22:421-478 (1988).

In some situations, a decreased oil content may be desired. A feature of the invention is that DAGAT activity can be reduced by gene silencing, antisense, ribozymes, cosuppression, or mutagenesis techniques, resulting in a decrease in oil content. Gene silencing techniques, such as that described in WO 98/36083 are useful. Antisense RNA has been used to inhibit plant target genes in a tissue-specific manner. See, for example, U.S. Patent Nos. 5,453,566, 5,356,799, and 5,530,192. Antisense nucleic acid constructs include a partial or a full-length coding sequence operably linked to at least one suitable regulatory sequence in antisense orientation.

Expression of DAGAT also can be inhibited by ribozyme molecules designed to cleave DAGAT mRNA transcripts. While various ribozymes that cleave mRNA at site-specific recognition sequences can be used to destroy DAGAT mRNAs, hammerhead ribozymes are particularly useful. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole requirement is that the target RNA contain a 5'-UG-3' nucleotide sequence. The construction and production of hammerhead ribozymes is

well known in the art. See, for example, U.S. Patent No. 5,254,678. Hammerhead ribozyme sequences can be embedded in a stable RNA such as a transfer RNA (tRNA) to increase cleavage efficiency in vivo. Perriman, R. et al., Proc. Natl. Acad. Sci. USA, 92(13):6175-6179 (1995); de Feyter, R. and Gaudron, J., Methods in  
5 Molecular Biology, Vol. 74, Chapter 43, "Expressing Ribozymes in Plants", Edited by Turner, P.C, Humana Press Inc., Totowa, NJ (1997). RNA endoribonucleases such as the one that occurs naturally in *Tetrahymena thermophila*, and which have been described extensively by Cech and collaborators also are useful. See, for example, U.S. Patent No. 4,987,071.

10 The phenomenon of co-suppression also has been used to inhibit plant target genes in a tissue-specific manner. Co-suppression of an endogenous gene using a full-length cDNA sequence as well as a partial cDNA sequence are known. See, for example, WO 94/11516, and U.S. Patent Nos. 5,451,514 and 5,283,124. Co-suppression of DAGAT activity in plants can be achieved by expressing, in the sense  
15 orientation, the entire or partial coding sequence of a DAGAT gene.

Mutagenesis can also be used to reduce acyltransferase activity in plants. Mutagenic agents can be used to induce random genetic mutations within a population of seeds or regenerable plant tissue. Suitable mutagenic agents include, for example, ethyl methyl sulfonate, methyl N-nitrosoguanidine, ethidium bromide, diepoxybutane,  
20 x-rays, UV rays, and other mutagens known in the art. The treated population, or a subsequent generation of that population, is screened for reduced oil content or reduced DAGAT activity that results from the mutation. Mutations can be in any portion of a gene, including the coding region, introns, and regulatory elements, that render the resulting gene product non-functional or with reduced activity. Suitable  
25 types of mutations include, for example, insertions or deletions of nucleotides, and transitions or transversions in the wild-type coding sequence. Such mutations can lead to deletion or insertion of amino acids, and conservative or non-conservative amino acid substitutions in the corresponding gene product.

#### *Characterization of Oils*

30 Techniques that are routinely practiced in the art can be used to extract, process, and analyze the oils produced by plants of the instant invention. Typically,

plant seeds are cooked, pressed, and extracted to produce crude oil, which is then degummed, refined, bleached, and deodorized. Generally, techniques for crushing seed are known in the art. For example, soybean seeds can be tempered by spraying them with water to raise the moisture content to, e.g., 8.5%, and flaked using a smooth roller with a gap setting of 0.23 to 0.27 mm. Depending on the type of seed, water may not be added prior to crushing. Application of heat deactivates enzymes, facilitates further cell rupturing, coalesces the oil droplets, and agglomerates protein particles, all of which facilitate the extraction process.

The majority of the seed oil can be released by passage through a screw press. Cakes expelled from the screw press are then solvent extracted, e.g., with hexane, using a heat traced column. Alternatively, crude oil produced by the pressing operation can be passed through a settling tank with a slotted wire drainage top to remove the solids that are expressed with the oil during the pressing operation. The clarified oil can be passed through a plate and frame filter to remove any remaining fine solid particles. If desired, the oil recovered from the extraction process can be combined with the clarified oil to produce a blended crude oil.

Once the solvent is stripped from the crude oil, the pressed and extracted portions are combined and subjected to normal oil processing procedures (i.e., degumming, caustic refining, bleaching, and deodorization). Degumming can be performed by addition of concentrated phosphoric acid to the crude oil to convert non-hydratable phosphatides to a hydratable form, and to chelate minor metals that are present. Gum is separated from the oil by centrifugation. The oil can be refined by addition of a sufficient amount of a sodium hydroxide solution to titrate all of the fatty acids and removing the soaps thus formed.

Deodorization can be performed by heating the oil to 500°F (260°C) under vacuum, and slowly introducing steam into the oil at a rate of about 0.1 ml/minute/100 ml of oil. After about 30 minutes of sparging, the oil is allowed to cool under vacuum. The oil is typically transferred to a glass container and flushed with argon before being stored under refrigeration. If the amount of oil is limited, the oil can be placed under vacuum, e.g., in a Parr reactor and heated to 500°F for the

same length of time that it would have been deodorized. This treatment improves the color of the oil and removes a majority of the volatile substances.

Oil content can be measured by NMR using AOCS Method AM 2-93 and AOCS Recommended Practice AK 4-95 or by near infra-red reflectance spectroscopy (NIR) using AOCS Method AK 3-94 and AOCS Procedure AM 1-92. Oil composition can be analyzed by extracting fatty acids from bulk seed samples (e.g., at least 10 seeds). Fatty acid TAGs in the seed are hydrolyzed and converted to fatty acid methyl esters. Percentages of fatty acids typically are designated on a weight basis and refer to the percentage of the fatty acid methyl ester in comparison with the total fatty acid methyl esters in the sample being analyzed. Seeds having an altered fatty acid composition may be identified by techniques known to the skilled artisan, e.g., gas-liquid chromatography (GLC) analysis of a bulked seed sample, a single seed or a single half-seed. Half-seed analysis is well known in the art to be useful because the viability of the embryo is maintained and thus those seeds having what appears to be a desired fatty acid profile may be planted to form the next generation. However, bulk seed analysis typically yields a more accurate representation of the fatty acid profile of a given genotype. Fatty acid composition can also be determined on larger samples, e.g., oil obtained by pilot plant or commercial scale refining, bleaching and deodorizing of endogenous oil in the seeds.

The following examples are to be construed as merely illustrative of how one skilled in the art can make and use the DAGAT gene fragments, and does not limit the scope of the invention described in the claims.

### EXAMPLES

Example 1 – Cloning of a Brassica Acyl Transferase: A mouse acyl CoA:diacylglycerol acyltransferase (DAGAT) gene was recently identified (Case et al., *Proc Natl Acad Sci USA* **95**:13018-13023, 1998). The deduced mouse DAGAT amino acid sequence was used to search for similar plant sequences in GenBank and dbEST databases. The mouse DAGAT protein sequence aligned with significant sequence identity and similarity to a putative *Arabidopsis thaliana* acyl-CoA:cholesterol acyltransferase (ACAT; GenBank Accession No. 3135276, locus ATAC003058). However, upon aligning the *A. thaliana* ACAT with the mouse

DAGAT and mouse ACAT (Case et al., *J Biol Chem* 273:26755-26764, 1998) sequences, it was found that the *A. thaliana* protein had greater homology to mouse DAGAT than to mouse ACAT. Thus, the type of acyltransferase encoded by the *A. thaliana* gene was unclear.

- 5           The genomic DNA sequence encoding the *A. thaliana* protein (GenBank Accession No. 3135276) was used to design two PCR primers for amplifying candidate DAGAT or ACAT genomic sequences from different sources. The 5' DAGAT-1 primer had the sequence  
caucaucaucauACTGCCATGGACAGGTGTGATTCTGCTTTTT TATCA (SEQ ID  
10 NO:1), and the 3' DAGAT-2 primer had the sequence  
cuacuacuacuaCTAGAGACAGGGCAATGTAGAAAGTATGTA (SEQ ID NO:2).  
Lowercase sequences were used for cloning into the pAMP1 vector (Gibco, BRL).

- PCR amplification using DAGAT-1 and DAGAT-2 primers was carried out as follows. Each 100µl PCR reaction mixture contained 50ng of genomic DNA, 200µM  
15 of each dNTP, 1X buffer B (Gibco BRL), 1 µM DAGAT-1 primer, 1 µM DAGAT-2 primer, 3mM magnesium sulfate, and 2µl Elongase enzyme (Gibco BRL). The reaction mixture was denatured at 94°C for 3 minutes, followed by 30 cycles of denaturation at 94°C for 1 minute, annealing at 50°C for 2 minutes, and extension at 72°C for 3 minutes. A final extension incubation was performed at 72°C for 10  
20 minutes after cycling. Based on the sequence of *A. thaliana* genomic DNA in GenBank Accession No. 3135276, a 1369 bp fragment was expected to be amplified from *Brassica*.

- An approximately 1.4 kb fragment was amplified from genomic DNA isolated from *Brassica napus* variety Westar under PCR conditions described above. The  
25 amplified DNA fragment was cloned into the pAMP1 vector (Gibco BRL) and partially sequenced from the 5' end using a T7 universal primer and from the 3' end using a Sp6 universal primer. Figure 1 contains a restriction map of the pAMP1 vector containing the putative DAGAT fragment, which has been designated pMB143. The partial sequence of the 5' end using the T7 primer was



ATGGACAGGTGTGATTCTGCTTTTTTATCAGGTGTCACGTTGATGCTCCTC  
 ACTTGCAATTGTGTGGCTGAAGTTGGTTTCTTACGCTCATACTAACTATGAC  
 ATAAGAACCCTAGCTAATTCATCTGATAAGGTAAAAGAAGTGATATAATA  
 TTGGTCACTTGCATTGTGTTTTACTATTTTGACCAGACACTGTTGAAAACCT  
 5 GTAGGCCAATCCTGAAGTCTCCTACTATGTTAGCTTGAAGAGCTTGGCGTA  
 TTTCATGCTTGCTCCACATTGTGTTATCAGGTAATCTGATGCGTCTTCTGC  
 TAATTGTATCATAATTATCTTTCACTTGCAAAAGTTTCTTGTCTAAAACCT  
 TCGCTCTTCGCTTTACCCAGCCGAGCTATCCACGTTCT (SEQ ID NO:3).

The partial sequence of the 3' end using the Sp6 primer was

10 ATCAATCTTGTCTTACTCAAAAATCATATTATGTTTACGTTANTAACCAAA  
 ATTCATGTACGCACTGTCTACCTTTGTCAGTATTGGAGAATGTGGAATATG  
 GTATGGTTCTCTTCTTGAACATCCCCTTCTTTTTTTATACAAAGCAGATTAA  
 GAAAAGCTTATTGAGATCTTGTTTTTTCTAATAGCCTGTTTATAAATGGAT  
 GGTTTCGACATGTATACTTTCCGTGCCTTCGCAGAAATATACCGAAAGTGA  
 15 GTGTAGTTAATTGCGATGATCGATATTTTTTCTGTGCTTCATAAATTTAAC  
 CCTCCACTCATTTTTTCCAGGTACCCGCTATTATCCTTGCTTTCTTAGTCT  
 CTGCAGTCTTTCATGAGGTATAATACATACTTTCTACATTGNCCCTGTCTC  
 (SEQ ID NO:4).

The sequence of the *B. napus* clone obtained using the T7 primer had an  
 20 overall identity of 76.5% with the corresponding region of the *A. thaliana* gene,  
 suggesting that the 1.4 kb cloned *B. napus* fragment encoded a protein that may be  
 related to the *A. thaliana* DAGAT or ACAT.

Plasmid pMB143 will be deposited with the American Type Culture  
 Collection (ATCC), 10801 University Blvd., Manassas, VA, 20110-2209.

25 Example 2 – Reduction of Oil Content in Brassica Plants Expressing  
Antisense DAGAT: The putative DAGAT genomic DNA cloned in pAMP1 was  
 excised with a SmaI/SnaBI digestion. The putative DAGAT fragment was purified  
 and inserted at the blunted, EcoRI site of pMB110 to generate pMB171. This  
 construct contains cruciferin promoter and cruciferin termination sequences. The  
 30 antisense orientation of the putative DAGAT insert in pMB171 was confirmed with  
 restriction analysis. The cruciferin promoter/putative DAGAT/cruciferin termination  
 cassette was released from pMB171 by partially digesting with XbaI and completely  
 digesting with XhoI, and cloned into the XhoI/XbaI sites of pMOG800 to generate  
 pMB170. Figure 2 provides a diagram of pMB170.

35 The construct pMB170 was used to transform *Agrobacterium* LBA4404. The  
 resulting *Agrobacterium* transformants were each co-cultivated separately with *B.*

*napus* variety Westar hypocotyls and cultured consecutively on incubation, selection (containing kanamycin) and regeneration media until green shoots were produced. Regenerated plantlets were transferred to the greenhouse and grown to maturity. Each T1 plant (N=56) was selfed and the resulting T2 seeds were harvested from each individual T1 plant.

Oil, protein, chlorophyll, glucosinolate, oleic acid (18:1), linoleic acid (18:2), and  $\alpha$ -linolenic acid (18:3) content were determined in the T2 seed samples. Oil content was measured by NIR using a Foss NIR Systems model 6500 Feed and Forage Analyzer (Foss North America, Eden Prairie, MN) calibrated according to manufacturer's recommendations. Canola seed samples, which represented wide ranges of the sample constituents listed above, were collected for calibration. Lab analysis results were determined using accepted methodology (*i.e.*, oil, AOCS Method Ak 3-94; moisture, AOCS Method Ai 2-75; fatty acid, AOCS Method CE 1e-91 and AOCS Method CE 2-66; chlorophyll, AOCS Method CC 13D-55; protein, AOCS Method BA 4e-93; and glucosinolates, AOCS Method Ak 1-92). Instrument response also was measured for each sample. A calibration equation was calculated for each constituent by means of chemometrics. These equations are combined into one computer file and are used for prediction of the constituents contained in unknown canola samples.

T2 seed samples containing unknown levels of the above constituents were prepared by removing foreign material from the sample. Cleaned whole seed was placed into the instrument sample cell and the cell was placed into the instrument sample assembly. Analysis was carried out according to manufacturer instructions and was based on AOCS Procedure Am 1-92. The results are predicted and reported as % constituent (% oil and protein are reported based on dry weight). Conversion from 'dry weight' basis to 'as is' basis for oil and protein can be calculated using the following formula:

$$\text{constituent (as is)} = \text{constituent (dry wt.)} \times [1 - (\% \text{ moisture}/100)].$$

The average dried oil content of the T2 seeds ( $39.62 \pm 1.95\%$ ,  $n=56$ ) was not significantly different from that of the control (Westar,  $39.92 \pm 1.63$ ,  $n=5$ ). There were ten T1 plants, however, that produced T2 seeds having a lower total oil content. In

these T2 seeds, the average oil content was  $36.63 \pm 0.64\%$ . Table 2 provides an analysis of eight of these 10 T2 seeds.

Approximately 10-20 T2 seeds from each of the ten selected plants were planted, and a nickel size portion of the leaf tissue was taken from plants about 2.5 weeks post-germination. Tissue samples were dried in a food dehydrator at  $135^{\circ}\text{C}$  for 8-16 hours. DNA was isolated using the Qiagen Dneasy96 Plant Kit, and resuspended in  $150\mu\text{l}$  of buffer. PCR amplification was performed in a  $20\mu\text{l}$  volume containing 1X PCR buffer containing  $1.5\text{mM}$   $\text{MgCl}_2$  (Qiagen PCR Core Kit),  $0.2\text{mM}$  dNTP,  $0.5$  units Taq polymerase (Qiagen),  $0.5\mu\text{M}$  cruciferen primer ( $5'$ -CTT TAT GGA TGA GCT TGA TTG AG- $3'$ , SEQ ID NO:5), and  $0.5\mu\text{M}$  acyltransferase primer ( $5'$ -CCG CTC TAG AGG GAT CCA AGC- $3'$ , SEQ ID NO:6),  $0.4\%$  sucrose, and  $0.008\%$  cresol. Amplification conditions included 30 cycles of denaturation at  $94^{\circ}\text{C}$  for 30 seconds, annealing at  $55^{\circ}\text{C}$  for 30 seconds, and extension at  $72^{\circ}\text{C}$  for 1 minute. PCR products were analyzed 1.2% agarose gel electrophoresis. As indicated in Table 1, the selected plants tested positive for the transgene.

**TABLE 2**  
**Analysis of T2 Seeds With Lower Oil Content**

SAMPLE	OIL	PROT	CHLOR	GLUC	C18:1	C18:2	C18:3	(# pos. plants/Total # plants tested)
Y350071	35.69	26.07	78.12	3.66	70.57	10.03	7.74	15/18
Y350070	36.35	24.87	78.57	2.60	72.02	8.18	7.34	11/18
Y350095	36.73	26.55	88.44	1.38	73.12	7.80	7.62	7/9
Y350079	36.79	27.96	22.75	2.96	66.67	14.97	7.53	17/18
Y350052	37.02	26.02	74.34	3.44	67.80	14.09	7.62	8/18
Y350077	37.04	25.70	70.80	2.91	71.30	9.16	8.07	15/18
Y350089	37.13	25.07	59.60	2.28	71.37	9.93	7.26	8/15
Y350078	37.66	25.64	43.00	2.35	74.75	7.48	6.94	11/18
Average	36.80	25.99	64.45	2.70	70.95	10.23	7.51	
Std Dev.	0.58	0.97	21.79	0.72	2.65	2.81	0.34	

**OTHER EMBODIMENTS**

It is to be understood that while the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of  
5 the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

## WHAT IS CLAIMED IS:

1. An isolated nucleic acid having at least 80% sequence identity to the nucleotide sequence of SEQ ID NO:3, or to a fragment of the nucleotide sequence of SEQ ID NO:3, said fragment being at least 15 nucleotides in length.
- 5 2. The nucleic acid of claim 1, said nucleic acid having at least 85% sequence identity to the nucleotide sequence of SEQ ID NO:3 or to said fragment.
3. The nucleic acid of claim 1, said nucleic acid having at least 90% sequence identity to the nucleotide sequence of SEQ ID NO:3 or to said fragment.
4. The nucleic acid of claim 1, said nucleic acid having at least 95%  
10 sequence identity to the nucleotide sequence of SEQ ID NO:3 or to said fragment.
5. An isolated nucleic acid having at least 80% sequence identity to the nucleotide sequence of SEQ ID NO:4, or to a fragment of the nucleotide sequence of SEQ ID NO:4, said fragment being at least 15 nucleotides in length.
6. The nucleic acid of claim 5, said nucleic acid having at least 85%  
15 sequence identity to the nucleotide sequence of SEQ ID NO:4 or to said fragment.
7. The nucleic acid of claim 5, said nucleic acid having at least 90% sequence identity to the nucleotide sequence of SEQ ID NO:4 or to said fragment.
8. The nucleic acid of claim 5, said nucleic acid having at least 95% sequence identity to the nucleotide sequence of SEQ ID NO:4 or to said fragment.
- 20 9. An isolated nucleic acid comprising a first and second region, said first region having at least 80% sequence identity to the nucleotide sequence of SEQ ID NO:3, said second region having at least 80% sequence identity to the nucleotide sequence of SEQ ID NO:4.
10. The nucleic acid of claim 9, wherein said nucleic acid encodes a  
25 diacylglycerol acyltransferase polypeptide.
11. The nucleic acid of claim 9, wherein said first and second regions are separated by approximately 600 nucleotides.
12. The nucleic acid of claim 9, wherein said nucleic acid is the insert of pMB143.

13. An expression vector comprising the nucleic acid of claim 9 operably linked to an expression control element.

14. The expression vector of claim 13, wherein said nucleic acid is operably linked in antisense orientation.

5 15. A transgenic plant comprising an exogenous nucleic acid encoding a diacylglycerol acyltransferase polypeptide operably linked to a regulatory element

16. The transgenic plant of claim 15, said nucleic acid comprising a first and second region, said first region having at least 80% sequence identity to the nucleotide sequence of SEQ ID NO:3, said second region having at least 80%  
10 sequence identity to the nucleotide sequence of SEQ ID NO:4.

17. The transgenic plant of claim 15, wherein said plant produces seeds that exhibit a statistically significantly altered oil content as compared to seeds produced by a corresponding plant lacking said nucleic acid encoding said diacylglycerol acyltransferase polypeptide.


15 18. The plant of claim 15, wherein said plant is a soybean plant.

19. The plant of claim 15, wherein said plant is a *Brassica* plant.

20. Seeds produced by the plant of claim 15.

21. Progeny of the plant of claim 15.

1/2

**FIG. 1**

SUBSTITUTE SHEET (RULE 26)

2/2




FIG. 2



## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/12316

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/54 C12N15/82 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EMBL, EPO-Internal, BIOSIS, WPI Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE GENBANK [Online] GI:3135250, 16 May 1998 (1998-05-16) ROUNSELEY, S.D., ET AL.: "Arabidopsis thaliana chromosome II BAC F27F23 genomic sequence complete sequence" XP002148213 see nts 91954-93322 F27F23.26 gene & DATABASE EMBL [Online] ACCESSION NO:AC003058, abstract & DATABASE GENPEPT [Online] GI:3135276 ACCESSION AAC16476 GENPEPT, 16 May 1998 (1998-05-16) ROUNSELEY, S.D., ET AL.: "putative acyl-CoA:cholesterol acyltransferase." abstract & DATABASE GENPEPT [Online] ACCESSION NO:T01294 GI:7488350, 30 April 1999 (1999-04-30) -/-	1,2,5,6, 9-11, 13-17, 20,21

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&amp;\* document member of the same patent family

Date of the actual completion of the international search

25 September 2000

Date of mailing of the international search report

20.10.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Maddox, A

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/12316

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>ROUNSLEY, S.D., ET AL.: "sterol O-acyltransferase homolog F27F23.26 - Arabidopsis thaliana." abstract</p> <p>---</p> <p>DATABASE EMBL [Online] ACCESSION NO:AC005917, 4 November 1998 (1998-11-04) LIN, X., ET AL.: "Arabidopsis thaliana chromosome II section 113 of 255 of the complete sequence. Sequence from clones F27F23, F3P11." XP002148214 see nts 22846-25865 At2g19450 -&amp; DATABASE GENBANK [Online] GI:4191771, 28 January 1999 (1999-01-28) LIN, X., ET AL.: "Arabidopsis thaliana chromosome II BAC F3P11 genomic sequence complete sequence." XP002148215 see nts 21175-24194 F3P11.5/F27F23.25 gene &amp; DATABASE STREMBL [Online] ACCESSION NO:Q9SLD2, abstract &amp; DATABASE GENPEPT [Online] ACCESSION AAD10144 GI:4191775, abstract</p>	1,2,5,6, 9-11, 13-17, 20,21
X	<p>---</p> <p>WO 98 55631 A (CALGENE LLC) 10 December 1998 (1998-12-10)</p>	15-21
Y	the whole document	15-21
X	<p>---</p> <p>HILLS M J ET AL: "DIACYLGLYCEROL ACYLTRANSFERASE - CLONING AND FUNCTIONAL EXPRESSION OF A CDNA FROM ARABIDOPSIS THALIANA" BIOCHEMICAL SOCIETY TRANSACTIONS,GB,COLCHESTER, ESSEX. 668TH MEETING OF THE BIOCHEMICAL SOCIETY; GLASGOW, SCOTLAND, UK; APRIL 7-9, 1999 , vol. 27, no. 3, 1999, page A124 XP000889768 ISSN: 0300-5127</p>	1-3,5-7, 9-11,13, 15-17, 20,21
Y	the whole document	15-21
X	<p>---</p> <p>DATABASE DBEST [Online] dbEST Id: 1277346, 19 September 1997 (1997-09-19) XP002148216 abstract &amp; DATABASE EMBL [Online] ACCESSION NO:AA042298, 5 September 1996 (1996-09-05) abstract</p> <p>---</p> <p>-/--</p>	1-3,5-7, 9-11,13, 15-17, 20,21

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/12316

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	KATAVIC ET AL: "Alteration of seed fatty acid composition by an ethyl methanesulfonate-induced mutation in Arabidopsis thaliana affecting diacylglycerol acyltransferase activity" PLANT PHYSIOLOGY, US, AMERICAN SOCIETY OF PLANT PHYSIOLOGISTS, ROCKVILLE, MD, no. 108, 1995, pages 399-409, XP002079129 ISSN: 0032-0889 the whole document	15,20,21
P,X	--- WO 99 63096 A (LASSNER MICHAEL W ; RUEZINSKY DIANE M (US); CALGENE LLC (US)) 9 December 1999 (1999-12-09)  see fig.1 and SEQ ID NO:1	1-7, 9-11,13, 15-17, 19-21
P,X	--- DATABASE EMBL [Online] ACCESSION NO:AJ131831, 10 June 1999 (1999-06-10) HILLS, M.J., ET AL.: "Arabidopsis thaliana mRNA for diacylglycerol 0-acyltransferase" XP002148217 the whole document -& HOBBS, D.H., ET A.: "Cloning of a cDNA encoding diacylglycerol acyltransferase from Arabidopsis thaliana and its functional expression" FEBS LETTERS, vol. 452, June 1999 (1999-06), pages 145-149, XP002122747	1-7, 9-11,13, 15-17, 20,21
P,X	--- DATABASE EMBL [Online] ACCESSION NO:AF155224, 30 June 1999 (1999-06-30) NYKIFORUK C.L., ET AL.: "Brassica napus putative diacylglycerol acyltransferase (DGAT2) mRNA, complete cds." XP002148218 the whole document -& CORY L. NYKIFORUK, ET AL.: "ISOLATION AND SEQUENCE ANALYSIS OF A NOVEL CDNA ENCODING A PUTATIVE DIACYLGLYCEROL ACYLTRANSFERASE FROM A MICROSPORE-DERIVED CELL SUSPENSION CULTURE OF BRASSICA NAPUS L. CV JET NEUF (ACCESSION NO. AF155224)." (PGR99-123) PLANT PHYSIOL. 120: 1207 , 1999, XP002148236 --- -/--	1-7, 9-13, 15-17, 19-21

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/12316

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>DATABASE EMBL [Online]  ACCESSION NO: AF164434,  26 July 1999 (1999-07-26)  NYKIFORUK C.L., ET AL.: "Brassica napus  putative diacylglycerol acyltransferase  (DGAT1) mRNA, complete cds."  XP002148219  the whole document  -&amp; CORY L. NYKIFORUK, ET AL.: "Isolation  and Characterization of a cDNA Encoding a  Second Putative Diacylglycerol  Acyltransferase from a Microspore-derived  Cell Suspension Culture of Brassica napus  L. cv Jet Neuf (Accession No. AF164434)."  PGR99-158) PLANT PHYSIOL. 121: 1057 ,  1999, XP002148237</p>	<p>1-7,  9-11,  15-17,  19-21</p>
P,X	<p>---  DATABASE EMBL [Online]  ACCESSION NO: AF251794,  17 April 2000 (2000-04-17)  BROWN, A.P., ET AL.: "Brassica napus  putative diacylglycerol acyltransferase  mRNA, complete cds."  XP002148220  the whole document</p>	<p>1-13,  15-17,  19-21</p>
P,X	<p>---  DATABASE BIOSIS [Online]  BIOSCIENCES INFORMATION SERVICE,  PHILADELPHIA, PA, US;  January 2000 (2000-01)  BOUVIER-NAVE PIERRETTE ET AL: "Expression  in yeast and tobacco of plant cDNAs  encoding acyl CoA:diacylglycerol  acyltransferase."  Database accession no. PREV2000000103940  XP002148221  abstract  &amp; EUROPEAN JOURNAL OF BIOCHEMISTRY,  vol. 267, no. 1, January 2000 (2000-01),  pages 85-96,  ISSN: 0014-2956  &amp; DATABASE EMBL [Online]  ACCESSION NO: AF051849,  23 December 1999 (1999-12-23)  abstract</p>	<p>1-7,15,  17,20,21</p>
P,X	<p>---  WO 00 01713 A (LARDIZABAL KATHRYN DENNIS  ;CALGENE LLC (US); HAWKINS DEBORAH (US);)  13 January 2000 (2000-01-13)  claims 1-48; examples 12,16  ---</p>	<p>15,17,  19-21</p>

-/--

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/12316

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 99 67403 A (UNIV CALIFORNIA) 29 December 1999 (1999-12-29) see SEQ ID NO:3 page 39, line 28 - line 32; claims 1-15 & WO 99 67268 A (UNIV CALIFORNIA) 29 December 1999 (1999-12-29) ---	15,17, 19-21
E	WO 00 36114 A (WEI YANGDOU ;ZOU JITAO (CA); JAKO COLETTE C (CA); TAYLOR DAVID C ( ) 22 June 2000 (2000-06-22)  the whole document ---	1-7, 9-11,13, 15-17, 20,21
E	WO 00 32756 A (DU PONT ;CAHOON EDGAR B (US); CAHOON REBECCA E (US); KINNEY ANTHON) 8 June 2000 (2000-06-08)  the whole document ---	1-7, 9-11,13, 15-17, 20,21
E	WO 00 32793 A (LARDIZABAL KATHRYN DENNIS ;METZ JAMES GEORGE (US); CALGENE LLC (US) 8 June 2000 (2000-06-08) the whole document ---	15,17-21
A	WESELAKE ET AL: "Differential display of mRNA from oil-forming cell suspension cultures of Brassica napus" PHYSIOLOGY, BIOCHEMISTRY AND MOLECULAR BIOLOGY OF PLANT LIPIDS,XX,XX, 1997, pages 357-359, XP002079130 the whole document ---	1-12
A	LITTLE ET AL: "Solubilization and characterization of diacylglycerol acyltransferase from microspore-derived cltures of oilseed rape" BIOCHEMICAL JOURNAL,GB,PORTLAND PRESS, LONDON, no. 304, 1994, pages 951-958, XP002079126 ISSN: 0264-6021 the whole document ---	1-12
A	WILSON R F ET AL: "RECENT DEVELOPMENTS IN THE MOLECULAR BIOCHEMISTRY AND GENETICS OF DIACYGLYCEROL ACYLTRANSFERASE FROM SOYBEAN" SEED OILS FOR THE FUTURE,US,AOCS PRESS, CHAMPAIGN, IL, 1992, pages 116-135, XP000884791 the whole document ---	1-12
	---	

-/--

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/12316

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>CASES S ET AL: "Identification of a gene encoding an acyl CoA: diacylglycerol acyltransferase, a key enzyme in triacylglycerol synthesis" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA,US,NATIONAL ACADEMY OF SCIENCE. WASHINGTON, vol. 95, no. 22, 27 October 1998 (1998-10-27), pages 13018-13023, XP002122745 ISSN: 0027-8424 the whole document</p>	1-12
A	<p>----- FRENTZEN M (REPRINT): "Acyltransferases from basic science to modified seed oils" FETT - LIPID,DE,WILEY-VCH VERLAG,WEINHEIM, vol. 100, no. 4/05, May 1998 (1998-05), pages 161-166, XP002122744 ISSN: 0931-5985 the whole document</p>	15-21
A	<p>----- WO 98 55632 A (CALGENE LLC) 10 December 1998 (1998-12-10) -----</p>	

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/12316

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9855631	A	10-12-1998	EP	1003882 A	31-05-2000
WO 9963096	A	09-12-1999	NONE		
WO 0001713	A	13-01-2000	NONE		
WO 9967403	A	29-12-1999	AU	4712499 A	10-01-2000
			AU	9037698 A	10-01-2000
			WO	9967268 A	29-12-1999
WO 0036114	A	22-06-2000	AU	1763800 A	03-07-2000
WO 0032756	A	08-06-2000	AU	1837600 A	19-06-2000
WO 0032793	A	08-06-2000	AU	2166400 A	19-06-2000
WO 9855632	A	10-12-1998	EP	0986647 A	22-03-2000

